6. $f(x) = 4x^3 + px^2 + 8x + q$

where p and q are constants.

Given that

- $(2x + 3)$ is a factor of $f(x)$
- $f(x)$ has a remainder of -5 when divided by $(x + 2)$

(a) (i) show that $p = 10$

(ii) find the value of q.

(5)

(b) Hence find the range of values of x for which $f(x)$ is decreasing.

(*Solutions based entirely on calculator technology are not acceptable.*)

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 9 marks)

7. **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

(i) Solve, for $0 \leqslant x < 2\pi$, the equation

$$3\sin x \tan x = 11 + \cos x$$

giving the answers in radians to 3 decimal places.

(5)

(ii) Given that

- $0 < \theta < 90°$
- $\cos\theta = \frac{1}{3}$

find, in simplest form, the exact value of $\tan\theta$

(2)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA