5.



Figure 2

Figure 2 shows a sketch of the curve defined by the parametric equations

$$x = t^2 + 2t \qquad y = \frac{2}{t(3-t)} \qquad a \leqslant t \leqslant b$$

where a and b are constants.

The ends of the curve lie on the line with equation $y = 1$

(a) Find the value of a and the value of b

(2)

The region R, shown shaded in Figure 2, is bounded by the curve and the line with equation $y = 1$

(b) Show that the area of region R is given by

$$M - k\int_a^b \frac{t+1}{t(3-t)}\,\mathrm{d}t$$

where M and k are constants to be found.

(5)

(c) (i) Write $\dfrac{t+1}{t(3-t)}$ in partial fractions.

(ii) Use algebraic integration to find the exact area of R, giving your answer in simplest form.

(6)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

Question 5 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 5 continued

(Total for Question 5 is 13 marks)

6. With respect to a fixed origin O, the line l_1 is given by the equation

$$\mathbf{r} = \mathbf{i} + 2\mathbf{j} + 5\mathbf{k} + \lambda(8\mathbf{i} - \mathbf{j} + 4\mathbf{k})$$

where λ is a scalar parameter.

The point A lies on l_1

Given that $|\overrightarrow{OA}| = 5\sqrt{10}$

(a) show that at A the parameter λ satisfies

$$81\lambda^2 + 52\lambda - 220 = 0$$

(3)

Hence

(b) (i) show that one possible position vector for A is $-15\mathbf{i} + 4\mathbf{j} - 3\mathbf{k}$

(ii) find the other possible position vector for A.

(3)

The line l_2 is parallel to l_1 and passes through O.

Given that

- $\overrightarrow{OA} = -15\mathbf{i} + 4\mathbf{j} - 3\mathbf{k}$
- point B lies on l_2 where $|\overrightarrow{OB}| = 4\sqrt{10}$

(c) find the area of triangle OAB, giving your answer to one decimal place.

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA