4. **In this question you must show all stages of your working.**
Solutions relying entirely on calculator technology are not acceptable.

(i) The volume, V, of a spherical balloon is increasing at a constant rate of $70\pi\,\text{cm}^3\,\text{s}^{-1}$

Find the rate of increase of the radius of the balloon, in $\text{cm}\,\text{s}^{-1}$, at the instant when the radius of the balloon is 5 cm.

[The volume V of a sphere of radius r is given by the formula $V = \frac{4}{3}\pi r^3$]

(4)

(ii) The depth of water in a cave is being monitored.

The rate of increase in the depth of water, h cm, at a particular point in the cave is modelled by the differential equation

$$\frac{\mathrm{d}h}{\mathrm{d}t} = \frac{k}{h^3}$$

where k is a constant and t hours is the time after monitoring began.

Given that

- initially the depth of water was 4 cm
- 5 hours after monitoring began, the depth of water was 6 cm
- T hours after monitoring began, the depth of water was 10 cm

solve the differential equation to find the value of T.

Give your answer to one decimal place.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

(Total for Question 4 is 10 marks)

5. (i) Find

$$\int x^2 e^{4x} dx$$

writing the answer in simplest form.

(4)

(ii) Use partial fractions and algebraic integration to show that

$$\int_4^7 \frac{2x+11}{(2x+1)(2-x)} dx = \ln k$$

where k is a fully simplified rational constant to be found.

(6)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA